

Appendix 5: Prevalence of resistance by country-level reported first-line antibiotic treatment for UTI in OECD (a) and non-OECD (b) countries

(a) OECD

Antibiotics	Reported <sup>a</sup> first-line UTI treatment				First-line UTI treatment not specified			
	Pooled prevalence (%)	Number of isolates tested	Number of reporting studies	I <sup>2</sup> (%)	Pooled prevalence (%)	Number of isolates tested	Number of reporting studies	I <sup>2</sup> (%)
<b>Ampicillin</b>	54.0 (44.5-63.4)	55,567	17 (6 countries)	69.6	52.7 (40.9-64.4)	14,617	8 (5 countries)	3.0
<b>Co-amoxiclav</b>	9.4 (5.9-12.9)	1971	7 (4 countries)	1.1	10.8 (7.7-14.0)	66,895	14 (5 countries)	0.0
<b>Co-trimoxazole</b>	31.5 (20.3-42.7)	53,634	23 (8 countries)	10.6	34.0 (24.9-44.1)	277	1 (1 country)	-
<b>Trimethoprim</b>	23.6 (17.9-30.3)	18,977	7 (5 countries)	11.5	-	0	0	-
<b>Nitrofurantoin</b>	2.4 (1.7-3.1)	51,486	7 (3 countries)	0.0	1.0 (0.2-1.8)	4801	14 (10 countries)	20.6
<b>Ciprofloxacin</b>	-	0	0	-	3.1 (0.8-5.4)	52,209	17 (9 countries)	0.0
<b>Ceftazidime<sup>b</sup></b>	3.1 (0.1-11.3)	22,962	5 (3 countries)	0.0	1.2 (0.4-2.0)	3020	5 (5 countries)	47.7

<sup>a</sup> Reported first-line treatment defined as either recommended as part of national/international guidelines or the antibiotic that would be given by a pharmacist when presenting with UTI symptoms.

<sup>b</sup> Ceftazidime reported as a marker for cephalosporin resistance, where any cephalosporin was reported as a first-line treatment.

## (b) Non-OECD

Antibiotics	Reported <sup>a</sup> first-line UTI treatment				First-line UTI treatment not specified			
	Pooled prevalence (%)	Number of isolates tested	Number of reporting studies	I <sup>2</sup> (%)	Pooled prevalence (%)	Number of isolates tested	Number of reporting studies	I <sup>2</sup> (%)
<b>Ampicillin</b>	84.2 (67.5-96.0)	1101	7 (6 countries)	0.0	77.9 (70.6-85.2)	1316	8 (5 countries)	39.6
<b>Co-amoxiclav</b>	-	0	0	-	60.3 (40.9-79.0)	1256	8 (8 countries)	69.7
<b>Co-trimoxazole</b>	76.2 (64.1-87.2)	2071	14 (7 countries)	7.3	55.6 (26.6-84.7)	661	4 (3 countries)	0.0
<b>Trimethoprim</b>	-	0	0	-	67.0 (64.4-72.4)	596	1 (1 country)	-
<b>Nitrofurantoin</b>	21.2 (17.0-24.8)	429	1 (1 country)	-	13.7 (7.6-19.7)	2782	17 (9 countries)	27.0
<b>Ciprofloxacin</b>	58.1 (51.5-64.7)	511	3 (2 countries)	0.0	15.8 (4.7-26.8)	1461	8 (5 countries)	26.3
<b>Ceftazidime<sup>b</sup></b>	22.3 (15.4-31.2)	113	1 (1 country)	-	24.2 (13.5-35.0)	1023	7 (4 countries)	63.1

<sup>a</sup> Reported first-line treatment defined as either recommended as part of national/international guidelines or the antibiotic that would be given by a pharmacist when presenting with UTI symptoms.

<sup>b</sup> Ceftazidime reported as a marker for cephalosporin resistance, where any cephalosporin was reported as a first-line treatment.